





## **AUTOLOGOUS CHONDROCYTE IMPLANTATION (ACI) OF PATELLOFEMORAL JOINT (PATELLA / TROCHLEA) WITH** TIBIAL TUBERCLE OSTEOTOMY AND MPFL RECONSTRUCTION

## PHYSICAL THERAPY PROTOCOL

Bryan M. Saltzman, M.D. OrthoCarolina Assistant Professor of Orthopaedic Surgery, Atrium Health Sports Medicine & Shoulder/Elbow 1915 Randolph Rd, Charlotte, NC 28207 704-323-3000 www.BryanSaltzmanMD.com

Patient Name: \_\_\_\_\_ Date of Surgery: \_\_\_\_\_

\_\_\_ Evaluate and Treat

\_\_\_\_ Provide patient with home program

Frequency: \_\_\_\_\_\_x/week x \_\_\_\_\_weeks

	WEIGHT BEARING	BRACE	ROM	EXERCISES
PHASE I	Heel-touch only	Locked in full extension at all times*	Gentle passive 0-45 <sup>°</sup>	Quad sets, patellar mobs, SLR, calf pumps at home
0-2 weeks		Off for hygiene and home exercise only		CPM at home





	CPM 0-30	





				OT IL RECEI OTIL RE
		2-4 weeks:	2-4 weeks:	2-6 weeks: Add side-lying hip and
	2-6 weeks:	Unlocked 0-45°	CPM 0-60°	core, advance quad set and stretching
PHASE II	Heel- touch only	4-6 weeks:	<b>4-6 weeks:</b> CPM 0-90°	<b>6-8 weeks</b> : Addition of heel raises, total gym (closed chain), gait
weeks	<b>6-8 weeks:</b> Advance 25% weekly until full	Unlocked 0-90°	Advance	normalization, eccentric quads, eccentric hamstrings
		Discontinue brace at 6	ROM as tolerated	Advance core, glutes and pelvic
		weeks	when non-WB	
PHASE				Progress closed chain activities
<b>III</b> 8-12	Full	None	Full	Advance hamstring work, proprioception/balance exercises; hip/core/glutes
weeks				Begin stationary bike at 10 wk
<b>PHASE</b> <b>IV</b> 12-24	Full	None	Full	Progress Phase III exercises and functional activities: walking lunges, planks, bridges, swiss ball, half-bosu exercises
weeks				Advance core/glutes and balance
PHASE V 6-12 months	Full	None	Full	Advance all activity w/o impact such as running, jumping, pivoting, sports until cleared by MD

\*Brace may be removed for sleeping after first post-operative visit (day 7-14)

By signing this referral, I certify that I have examined this patient and physical therapy is medically necessary. This patient \_\_\_\_\_ would \_\_\_\_\_ would not benefit from social services.

Date:\_\_\_\_\_

Bryan M. Saltzman, MD