





ANTERIOR CRUCIATE LIGAMENT (ACL) AND **POSTEROLATERAL CORNER (PLC) RECONSTRUCTION**

PHYSICAL THERAPY PROTOCOL

Bryan M. Saltzman, M.D.

OrthoCarolina Assistant Professor of Orthopaedic Surgery, Atrium Health Sports Medicine & Shoulder/Elbow 1915 Randolph Rd, Charlotte, NC 28207 704-323-3000 www.BryanSaltzmanMD.com

Patient Name: _____ Date of Surgery: _____

Evaluate and Treat

____ Provide patient with home program

Frequency: ______x/week x _____weeks

	WEIGHT BEARING	BRACE	ROM	EXERCISES
PHASE I	Heel touch WB in brace *	ambulation and	0-2 weeks:	Quad sets, patellar mobs, gastroc/soleus stretch SLR w/ brace in full extension until quad strength prevents extension lag
0-6 weeks		2-6 WEEKS:	2-6 weeks : Advance slowly 0-90	Side-lying hip/core Hamstrings avoidance until 6 wks post- op
PHASE	Advance 25%	Discontinue at 6	Full	Begin toe raises, closed chain quads,





				OTTICCUONITIC
II 6-12 weeks	weekly until full by 8 wks	wks if no extension lag		balance exercises, hamstring curls, stationary bike, step-ups, front and side planks; advance hip/core
PHASE III 12-16	Full	None	Full	Advance closed chain strengthening Progress proprioception activities Begin stairmaster, elliptical and running
weeks PHASE				straight ahead at 12 weeks 16 wks : Begin jumping
IV 16-24 weeks	Full	None	Full	20 wks : Advance to sprinting, backward running, cutting/pivoting/changing direction, initiate plyometric program and sport-specific drills
PHASE V > 6 months	Full	None	Full and pain- free	Gradual return to sports participation after completion of FSA*** Maintenance program based on FSA

*Modified with concomitantly performed meniscus repair/transplantation or articular cartilage procedure **Brace may be removed for sleeping after first post-operative visit (day 7-10)

***Completion of FSA (Functional Sports Assessment) not mandatory, but recommended at 22-24 wks post-op for competitive athletes returning to play after rehab

By signing this referral, I certify that I have examined this patient and physical therapy is medically necessary. This patient _____ would _____ would not benefit from social services.

Bryan M. Saltzman, MD