



POSTERIOR INSTABILITY REPAIR (LABRAL STABILIZATION)

PHYSICAL THERAPY PROTOCOL

Bryan M. Saltzman, M.D.

OrthoCarolina

Assistant Professor of Orthopaedic Surgery, Atrium Health

Sports Medicine & Shoulder/Elbow

1915 Randolph Rd, Charlotte, NC 28207

704-323-3000

www.BryanSaltzmanMD.com

Patient: _____

Date of Surgery: _____

Procedure: Right / Left Shoulder Posterior Stabilization

___ Evaluate and Treat

___ Provide patient with home program

Frequency: _____x/week x _____weeks

Phase I (0-6 wks): *Period of protection: In general, sling should be worn at all times during this phase (except for hygiene and PT). Motion and strengthening exercises are performed within strict motion limits.*

Weeks 0-3: No formal PT.

- Shoulder immobilizer should be worn at all times (except for hygiene and pendulums).



- Home exercises only (pendulums, elbow + wrist ROM, grip strengthening).

Weeks 3-6: Begin formal PT (2-3 x/wk).

- Sling at all times (except for hygiene and PT).
- ROM: Restrict motion to 90 deg FF / IR to the stomach / ER as tolerated with arm at side. **No cross-body adduction.**
 - Progress PROM → AAROM → AROM as tolerated within the above limits
 - Heat before, ice after.
- Strengthening: Cuff/periscapular/deltoid isometrics in sling, within above motion limits.

Phase II (6-12 wks): Advance active motion and strengthening.

- D/C sling if cleared by MD
- ROM: Progress active ROM to within 20 degrees of opposite side; avoid aggressive passive stretching in forward flexion, cross-body adduction and IR.
- Strengthening:
 - Progress cuff/deltoid and periscapular strengthening: isometrics → bands → light weights (1-5lbs) w/8-12 reps x 2-3 sets for cuff, deltoid, scapular stabilizers (*Only do this 3x/wk to avoid cuff tendonitis*)
 - Modalities as per PT discretion

Phase III (3-12 months): Progress to sport/occupation-specific rehab.

- ROM: Passive stretching at end ranges if full motion not yet achieved, as tolerated.
- Strengthening/Activities:
 - Continue bands/light weights as above, 3x/wk.
 - Begin eccentrically resisted motions, plyometrics (*weighted ball toss*), proprioception (*body blade*) and progress to sport-specific/job-specific exercises by 4 months.
 - **Throwers**:
 - @ 4.5 months, may begin light-tossing if full-strength and motion.
 - @ 6 months throw from the pitcher's mound and/or return to collision sports (hockey, football, etc.).
 - **Work**:



- Overhead work without lifting is usually possible @ 4.5-6 months.
- Can resume heavy labor once full-strength achieved (usually 6-9 months).

By signing this referral, I certify that I have examined this patient and physical therapy is medically necessary. This patient ___ would ___ would not benefit from social services.

Date: _____

Bryan M. Saltzman, MD